



Dorotea



seasonality

Late primocane variety,
30 days later than Enrosadira®.
Mid-Late florican variety,
10 days later than Enrosadira®.



climatic areas

Versatile variety. It shows its best
in areas with at least 800 chill hours, but
it shows nice yield also
with lower amount of chill.



plant structure and agronomic description

Vigorous and erect plant, with low
amount of spines. High amount of
flowers. Medium-short lateral shoot,
good in flower pollination. **Auto-fertile
variety**, but the presence of pollinator
insects is always helpful.



yield

High yield variety, suitable
for high density cultivation systems.
Quite easy to pick.



berry description

Medium conical shape, big size
and good firmness.
Exceptional taste.



RIPENING PERIOD

	May				June				July				August				September				October				
	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	
ENROSADIRA®				■	■	■	■							■	■	■	■	■	■	■	■				
HALLEY				■	■	■	■	■						■	■	■	■	■	■	■	■	■			
CASTION®					■	■	■	■	■						■	■	■	■	■	■	■	■	■		
DOROTEA						■	■	■	■	■						■	■	■	■	■	■	■	■	■	■
AURORA®						■	■	■	■	■	■						■	■	■	■	■	■	■	■	■

Data collected from Vivai Molari in Trento (2018/2019)

VARIETY	SEASONALITY	CHILL REQUIREMENT	BERRY WEIGHT (g)	BERRY COLOR*
ENROSADIRA®	Early	Mid - High	6	light red
HALLEY	Early	Mid - High	5,3	red
CASTION®	Early - Mid	Mid - High	6,6	shiny red
DOROTEA	Mid - Late	Mid - High	7,1	red
AURORA®	Late	Mid - High	5,7	pink red
POLKA	Early	High	6,6	red
AUTUMN BLISS	Early	High	4	red

*BERRY COLOR at harvest day