



Halley



seasonality

Early ripening primocane variety, 5 days later than Enrosadira®.

Early ripening florican variety, same as Enrosadira®.



climatic areas

Very versatile variety, adapted from Continental to Low Chill Climates.

Showing its best potential in areas with at least 400 chill hours.



plant structure and agronomic description

Medium length of the canes, thick canes, completely thornless. Medium length of the lateral shoots. Good flower pollination. **Auto-fertile variety,** but the presence of pollinator insects is always helpful.



yield

High yield variety, suitable for high density cultivation systems. Really easy to pick and to cultivate, being completely thornless.



berry description

Conical shape, medium to large size and very firm berries, light red color, sweet, **low acid** and **very aromatic berries.**



RIPENING PERIOD

	May				June				July				August				September				October			
	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4
ENROSADIRA®																								
HALLEY																								
CASTION®																								
DOROTEA																								
AURORA®																								

Data collected from Vivai Molari in Trento (2018/2019)

VARIETY	SEASONALITY	CHILL REQUIREMENT	BERRY WEIGHT (g)	BERRY COLOR*
ENROSADIRA®	Early	Mid - High	6	light red
HALLEY	Early	Mid - High	5,3	red
CASTION®	Early - Mid	Mid - High	6,6	shiny red
DOROTEA	Mid - Late	Mid - High	7,1	red
AURORA®	Late	Mid - High	5,7	pink red
POLKA	Early	High	6,6	red
AUTUMN BLISS	Early	High	4	red

*BERRY COLOR at harvest day